

Craven[®]

Craven is a non-volatile, fast acting herbicide desiccant that is registered for use on pulse, oilseed, and legume forage seed crops.

Registered Uses:

- Desiccation of pulse, oilseed, and legume forage seed crops.
- Weed control in vegetable and field crops.
- Control of Corn Spurry in oats.
- Potato Vine killing.
- Weed control in non-crop land.

Application Tips:

Craven application should be considered when a crop starts to change color. For a summary of ground rates and application volumes, see Table 1 on page 2. For complete details regarding rates, application volumes, and crop specific growth stages, please refer to the Craven label.

- Always use recommended water volume as complete coverage is essential.
- Craven is rapidly absorbed by plants and effectiveness is not reduced by rain shortly after treatment. Effectiveness of the treatment may be enhanced when application is made on cloudy days or prior to periods of darkness.
- Be aware that Craven accelerates the natural process of crop dry down. In case of adverse weather conditions such as heavy rain, hail or strong wind, the resulting damage to your crop may be enhanced. Take note that certain crops are more fragile than others.
- Use HiActivate, a non-ionic surfactant at a rate recommended on the label.

Harvesting Tips:

The use of Craven facilitates direct combining of many field crops such as lentils, peas, canola*, mustard or legumes.

- Farmers who wish to swath desiccated crops should wait until the crop has dried down sufficiently to allow the crop to be picked up and threshed immediately after swathing.
- Delaying threshing after swathing desiccated crops will increase shattering and seed loss.
- For most crops, harvest can normally commence within 4-10 days after desiccation. Adverse weather conditions can delay commencement of harvest beyond 10 days after application. When those conditions prevail after Craven desiccation, commence harvest when plant material is dry and seed moisture level allows efficient harvesting.
- To minimize seed loss and to maintain seed quality, harvest of desiccated crops should commence as soon as seed moisture reaches the level for normal harvest.

*The product can cause shattering losses in canola. It should only be used on B.napus canola to assist in the harvest of a severely lodged crop.



REGION

Western Canada



USE RATE

See adjacent table



APPLICATION GUIDE

See adjacent table



PACKAGING

115 L



ACTIVE INGREDIENTS

Diquat Dibromide 240g/L



HERBICIDE GROUP

Group 22



Craven®

Table 1 – Summary of ground application rates and application volumes:

For a complete details regarding ground and aerial rates, application volumes, and crop specific growth stages, please refer to the Craven label.

FIELD CROPS	RATE (L/ha)	WATER VOLUME (L/ha)	AREA TREATED (ha) / 115L
Beans - White & Red Kidney, Soybeans and Adzuki Beans	1.25 - 1.7	225-550	67.6 - 92
Canola*	1.25 - 1.7	225-550	67.6 - 92
Chickpeas	1.25 - 1.7	225-550	67.6 - 92
Flax (including low linolenic acid varieties)	1.25 - 1.7	225-550	67.6 - 92
Legumes (alfalfa, birdsfoot trefoil, red clover and white clover) Seed Crops	1.7 - 2.7	225-550	42.6 - 67.6
Lentils	1.25 - 1.7	225-550	67.6 - 92
Mustard (condiment type only)	1.25 - 1.7	225-550	67.6 - 92
Oats - Corn Spurry Control	0.9 - 1.25	225-335	92 - 127.8
Peas - Field or Dry	1.25 - 1.7	225-550	67.6 - 92
Potato - Vine Killing	1.25 - 3.5	550-1100	32.9 - 92
Sweet White Lupins	2.3	225-550	50
Sunflowers	1.25 - 1.7	225-550	67.6 - 92
VEGETABLES AND FIELD CROPS	RATE (L/ha)	APPLICATION VOLUME (L/ha)	AREA TREATED (ha) / 115L
Stale Seedbed	2.3 - 4.6	at least 300	25 - 50
Inter-row Directed Weeding	2.3 - 4.6	900-1100	25 - 50
FRUIT	RATE (L/ha)	APPLICATION VOLUME (L/ha)	AREA TREATED (ha) / 115L
Perennial Grass Suppression Under Apple Trees	4.6	225-675	25
NON-CROP LAND	RATE (L/ha)	APPLICATION VOLUME (L/ha)	AREA TREATED (ha) / 115L
Weed Control in Non-crop Land	2.3 - 4.6	550-1100	25 - 50

*The product can cause shattering losses in canola. It should only be used on B.napus canola to assist in the harvest of a severely lodged crop.

