

Antler®

For use in canola, pulses, and other broadleaf crops, Antler is a post-emergent herbicide for control of grassy weeds.

Antler is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within one to three weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and complete control three to five weeks after application solution.

FEATURES AND BENEFITS

- Can be tank-mixed with a wide range of broadleaf tank mix options, including Liberty®, to provide broad spectrum weed control.
- Can be used across a wide range of broadleaved crops.
- Wide window of application timing.
- Rainfast in 1 hour.

For use on (including but not limited to*)

Canola | Chickpeas | Field Peas | Flax | Lentils | Mustard | Potatoes | Soybeans
*see label for full listing of registered crops

Application Timing

Most crops are tolerant at all stages, so target applications at the optimal weed stage. See label for more information.

Weeds	Leaf Stage	Rate
Wild Oats, Volunteer Oats, Green & Yellow Foxtail, Volunteer Cereals (wheat, barley, oats)	2-4	50 mL/ac
Barnyard grass, Fall panicum, Proso millet, Volunteer corn, Volunteer canarygrass, Witch grass	2-6	50 mL/ac
Barnyard grass, Crabgrass (smooth, large), Fall panicum, Foxtail (green, yellow), Persian darnel, Proso millet, Quack grass suppression, Volunteer canarygrass, Volunteer cereals (wheat, oats, barley), Volunteer corn, Wild oats, Witch grass	2-6	80 mL/ac
Quackgrass control	2-4	150 mL/ac



REGION

All



USE RATE

See adjacent table



APPLICATION GUIDE

See adjacent table



PACKAGING

3 L + 9 L



ACTIVE INGREDIENTS

Clethodim 240 g/L



HERBICIDE GROUP

Group 1