

Marshall®

Controls a wide range of grassy weeds in pulse, oilseed and soybean crops

Marshall herbicide contains quizalofop P-ethyl, a post-emergent group 1 herbicide for control of annual and perennial grass in broadleaf crops across Canada.

FEATURES AND BENEFITS

- Consistent control of winter annual and perennial grasses like foxtail barley, bromes, volunteer cereals, and wild oats
- Effective control of glyphosate tolerant volunteer corn in glyphosate tolerant soybeans
- Excellent crop safety on all labeled crops
- Rainfast in 1 hour
- Tank mix compatible with many other broadleaf herbicides

For use on (including but not limited to*)

Canola | Chickpeas | Field Peas | Flax | Lentils | Seed Alfalfa | Soybeans
*see label for full listing of registered crop

Adjuvant Sold Separate

Marshall must be tank-mixed with an adjuvant to achieve optimal weed control. Marshall may be tank-mixed with various adjuvants, including crop oil concentrates (COC), methylated seed oils (MSO), and non-ionic surfactants (NIS). In optimal growing conditions where weeds are small, an NIS like HiActivate can be effective. In less optimal growing conditions, such as drought conditions, or when weeds are large in size, an MSO like Destination™ will provide superior performance.

When tankmixing with other herbicides select an adjuvant recommended for use with both products.

Application Information

Adjuvant	RATE
Crop Oil Concentrate	0.50 - 1.0% v/v
Methylated Seed Oil	0.50 - 1.0% v/v
Non Ionic Surfactant	0.25 - 0.5% v/v



REGION
All



USE RATE
Water Volume
Ground : 10 US gal/ac
Aerial: >3 US gal/ac

PRODUCT RATE:
0.15 - 0.3 L/ac
Refer to label for crop specific recommendations

SURFACTANT
MSO 0.50 - 1.0% v/v
NIS 0.25 - 0.50% v/v



APPLICATION GUIDE
See adjacent table



PACKAGING
2 x 10 - L



ACTIVE INGREDIENTS
Quizalofop - Group 1

Always read and follow label directions.
© 2020 WinField United. Destination™, HiActivate®, Marshall®, and WinField® are registered trademarks of WinField United.
All other trademarks are property of their respective owners.



Application Rates

Weeds	Leaf Stage	Rate
Green Foxtail, Volunteer Barley, Corn, Oats, and Wheat	2 leaf to early tillering	0.15 L/acre (67 acres/jug)
Wild Oats	1-5 leaf without tillers	
Barnyard Grass	2 leaf to early tillering	0.2 L/acre (50 acres/jug)
Wild Oats	*1-5 leaf to early tillering (up to 2 tillers)	
Fall Panicum, Proso Millet, and Yellow Foxtail,	2 leaf to early tillering	
Foxtail Barley	3 leaf to 4 leaf +tillers	
Downey and Japanese Brome	2-5 leaf	0.3 L/ac (33 acres/jug)
Weeds Controlled: Quackgrass	2-6 leaf	

*best results on volunteer/wild oats if application is made before tillering begins

NOTE: Use the higher rate of Crop Oil Concentrate when wild oats or quackgrass are the main target weeds, or under conditions not conducive to good growth.

